



ACC.15

TCT@ACC-12 | innovation in intervention

A566
JACC March 17, 2015
Volume 65, Issue 10S

Congenital Heart Disease

ELECTIVE BALLOON PULMONARY VALVULOPLASTY IN THE NEWBORN WITH TETRALOGY OF FALLOT

Poster Contributions

Poster Hall B1

Monday, March 16, 2015, 9:45 a.m.-10:30 a.m.

Session Title: Pediatric Congenital Heart Disease: Interventional

Abstract Category: 11. Congenital Heart Disease: Pediatric

Presentation Number: 1256-323

Authors: *Mohamed Leye, Nagib Dahdah, Joaquim Miro, CHU Sainte Justine, Montréal, Canada*

Background: In tetralogy of Fallot (TOF), the risk of cyanotic spells can lead to emergency systemic-pulmonary shunt. The development of the pulmonary artery and the onset of cyanotic spells are conditioned by the volume of the trans-valvular flow.

Methods: We conducted a retrospective study to determine whether elective primary pulmonary valvuloplasty (EPPV) in the neonate 1) allows a better development of the pulmonary artery; 2) reduces the risk of cyanotic spells or the use of emergency surgery; 3) increase the possibilities of conservative surgery of the pulmonary valve. TOF who received EPPV within the first month of live were compared to an observational group (OBS).

Results: There were 48 patients (19 EPPV and 29 OBS) with a trend for lower birth O₂-saturation in EPPV ($89\pm 5.7\%$ vs $94\pm 5.9\%$; $p=0.07$). Pulmonary valve z-score was comparable (-2.17 ± 1.43 vs -1.87 ± 1.54 ; $p=0.513$) as well as Nakata index (120.9 ± 41.6 vs 147.4 ± 50.1 mm²/m²; $p=0.62$). Pulmonary valve z-score remained comparable at surgery (-1.63 ± 1.83 vs -1.09 ± 1.75 ; $p=0.603$) and the Nakata index increased equally by 45.1 ± 97.8 vs 18.6 ± 56.8 ; $p=0.293$). Cyanotic spells and palliative shunts occurred equally in both groups (3 vs 2; $p=0.372$, and 1 vs 1; $p=1.0$, respectively). The need for trans-annular patch was higher in EPPV however (78.9% vs 34.5% ; $p=0.03$).

Conclusion: Despite appropriate growth of the pulmonary arteries following neonatal EPPV in TOF, there was no significant decrease of cyanotic spells, or the need for transannular patch.